

Current-like variables in massive and massless integrable models¹

L.Faddeev

*St.Petersburg Branch of the Steklov Mathematical Institute
Fontanka 27, St.Petersburg 191011, Russia*

and

*Research Institute for Theoretical Physics
P.O. Box 9 (Siltavuorenpenger 20C), SF-00014 University of Helsinki, Finland*

1 Introduction

In these lectures I shall deal with the representative examples of integrable models of quantum field theory in $1 + 1$ dimensional space-time. The term "integrable" first used in this domain in [1], means that the equations of motion have enough conservation laws to be exactly soluble. In quantum case the spectrum and eigenvectors of the Hamiltonian should be calculable. Twenty years of development showed that it is indeed the case and the famous Bethe Ansatz [2] found its reincarnation as a main tool to make it possible [3]. A parallel development was done in the 2-dimensional lattice classical statistical mechanics [4].

While the latter development was highly appreciated by the physics community, the former was considered rather peripheral because of its main limitation - space-time being two-dimensional. String theory [5] changed the situation, when the two-dimensional world-sheet acquired a bona fide physical interpretation. However it were the conformal (massless) examples of field theoretical models [6], which become prominent in this connection. The known (massive) examples of integrable models were resurrected as a perturbation (deformation) of integrable models [7] and exist now in the minds of young generation in this guise.

For me as a specialist working on integrable models for more than twenty years the opposite picture is more natural : the conformal field theory models are contractions of integrable models. One can construct and classify integrable models (at least classically [8]) without any reference to their conformal contractions. The latter are to appear afterwards as particular massless limits.

In these lectures I shall discuss some technique and results pertinent to the ideology, underlined above. It is natural to treat the massive and massless models on some common ground. The massive models have divergences and must be regularized. One of the nice features of their theory is that one can devise the lattice formulation where the integrability remains intact. Thus we should deal with the lattice models. On the other hand, the massless models, having chiral excitations (left and (or)

¹Lectures delivered at the International School of Physics "Enrico Fermi", held in Villa Monastero, Varenna, Italy, from 28 June to 8 July 1994

right movers), use the current-like dynamical variables. Thus, one is to devise the way of using the current-like variables for the massive models instead of the traditional canonical pairs. Both goals will be accomplished in these lectures. Finally, we shall work with finite volume space, which is natural for the conformal field models.

2 Dynamical variables

Lattice and finite volume regularizations lead to consideration of quantum mechanical system with finite number of degrees of freedom. As basic generators of the algebra of observables we shall take, following [9], [10], the set of dynamical variables w_n , $n = 1, \dots, N$, subject to the commutation relations

$$w_n w_{n+1} = q^2 w_{n+1} w_n,$$

$$w_n w_m = w_m w_n, \quad |n - m| \geq 2,$$

and the periodicity condition

$$w_{n+N} = w_n.$$

Here q is a complex number, playing the role of coupling constant. In most of applications q takes values on the circle

$$|q| = 1,$$

or

$$q = e^{i\gamma\hbar},$$

where γ is real and we introduced explicitly the Planck constant \hbar . This requirement is compatible with the unitarity of w_n :

$$w_n^* = w_n^{-1}.$$

However, from time to time it will be convenient to relax this condition.

Algebra \mathcal{A} , generated by w_n , has one central element

$$C = w_1 w_2 \dots w_N$$

for N odd and two central elements

$$C_1 = w_1 w_3 \dots w_{N-1}, \quad C_2 = w_2 w_4 \dots w_N$$

for N even. Thus we have $\frac{N-1}{2}$ degrees of freedom for N odd and $\frac{N}{2} - 1$ for N even. In that follows we shall consider N to be even ($N = 2M$), so that we shall deal with $M - 1$ degrees of freedom.

We can realize the generators w_n in the form

$$w_n = e^{ip_n},$$

if the p_n have the following commutation relations

$$[p_n, p_m] = 2i\hbar\gamma(\delta_{n+1,m} - \delta_{n,m+1}).$$

However, the substitution

$$w_n \rightarrow p_n = \frac{1}{i} \ln w_n$$

is not harmless; in particular, variables p_n belong not to the algebra \mathcal{A} , but rather to its extension. For evident reasons we shall refer to algebra \mathcal{A} , generated by w_n , as a compact one and to larger algebra $\hat{\mathcal{A}}$, generated by p_n , as a noncompact.

The algebra $\hat{\mathcal{A}}$ is much larger than \mathcal{A} . However, one can use w -like operators to describe it. For this goal we shall introduce the variables :

$$\hat{w}_n = e^{i\frac{\pi}{\gamma} p_n} = w_n^{\frac{\pi}{\gamma}}.$$

It is evident that the variables w_n and \hat{w}_n commute

$$\hat{w}_n w_m = w_m \hat{w}_n.$$

It is a less trivial statement, that w and \hat{w} together generate all $\hat{\mathcal{A}}$ for any q not being a root of unity. Algebra \mathcal{A} then is a factor inside of $\hat{\mathcal{A}}$.

To make these statements more exact one has to introduce some topology, which is standard (at least for $|q| = 1$), and use the language of v-Neumann algebras. I do not have time to do it in these lectures. One thing is remarkable and worth to be mentioned explicitly. The algebra \mathcal{A} has a trace at least for $C_1 = C_2 = 1$! If a generic element is given by

$$a = \sum a_{i_1 \dots i_{N-2}} w_1^{i_1} \dots w_{N-2}^{i_{N-2}},$$

(we do not use the dependent variables w_{N-1}, w_{N-2}), then

$$\text{tr } a = a_{0,0 \dots 0}.$$

Using the commutation relations one can show that the main property of trace

$$\text{tr}(ab) = \text{tr}(ba)$$

holds. The trace of unity is equal to 1, so we have here the example of a finite factor and in general factor II_1 .

The complimentary variables \hat{w}_n have the same current-like commutation relations

$$w_{n+1} \hat{w}_n = \hat{q}^2 \hat{w}_n w_{n+1},$$

where

$$\hat{q} = q^{-\frac{\pi^2}{\gamma^2}}.$$

The role of the modular map

$$\frac{\gamma}{\pi} \rightarrow -\frac{\pi}{\gamma}$$

entering here will be commented on later.

The described realisation of w via p is appropriate for the massless case. Indeed, naive classical ($\hbar \rightarrow 0$) and continuous limit, realized via rescaling

$$p_n = 2\Delta p(x), \quad x = n\Delta,$$

with $\Delta \rightarrow 0$ leads to the Poisson brackets

$$\{p(x), p(y)\} = \gamma \delta'(x - y)$$

typical for current-like variables.

However, the same algebra can be realized via canonical pairs (ϕ, π) with the commutation relations

$$[\phi_n, \pi_m] = -i\hbar \gamma \delta_{nm}, \quad n, m = 1, \dots, M$$


for example as

$$w_{2n} = e^{2i\gamma\phi_n}, \quad w_{2n+1} = e^{i(\pi_n - \pi_{n-1})}.$$


Here the continuous limit exists for

$$\phi(x) = \phi_n, \quad \Delta\pi(x) = \pi_n, \quad x = n\Delta.$$

The picture for space is



for the first case and



for the second one. The first is to be used in massless case, while the second one for the massive case.

Together with the current-like variables one can use also the vertex-like ones. In the continuous case they are defined by

$$p(x) = \psi'(x).$$

On a lattice massless and massive cases are to be treated separately. We shall need only the second case. The analogue of space derivative is the ansatz ;

$$w_n = \frac{\psi_{n+1}}{\psi_{n-1}}.$$

The inverse map $w \rightarrow \psi$ involves the product

$$\psi_n = \prod_{k \leq n} w_k,$$

where k and n have an opposite parity.

The last formula needs some clarification with respect to periodicity. Instead we shall introduce the algebra with generators ψ_n independently. The whole algebra (denoted as \mathcal{B}) is generated by ψ_n , $n = 1, \dots, 2M$, C_1 and C_2 with relations

$$\psi_n \psi_m = q \psi_m \psi_n$$

for $(n - m)$ odd; the quasiperiodicity holds as

$$\psi_{2n+2M} = C_1 \psi_{2n},$$

$$\psi_{2n+1+2M} = C_2 \psi_{2n+1},$$

finally

$$\psi_{odd} C_1 = q^2 C_1 \psi_{odd},$$

$$\psi_{even} C_2 = q^2 C_2 \psi_{even},$$

and ψ_{even} (ψ_{odd}) commute with C_1 (C_2). This algebra has no central elements, so the number of degrees of freedom is equal to $M + 1$. It may be reduced to the algebra \mathcal{A} , generated by w_n , if we introduced two constraints, fixing the values of (commuting) generators C_1 and C_2 .

One can construct almost canonical basis, using some mixture of w and ψ generators. Indeed, pairs (w_{even}, ψ_{even}) or (w_{odd}, ψ_{odd}) are Weyl pairs, i.e.

$$w_{2n} \psi_{2n} = q^2 \psi_{2n} w_{2n}, \quad n = 1, \dots, M,$$

$$w_{2n} \psi_{2m} = \psi_{2m} w_{2n}, \quad n \neq m$$

and give us the system with M degrees of freedom. Here the variable C_1 enters via the quasiperiodic conditions for ψ_{even} . The same is true for $(w_{odd}, \psi_{odd}, C_2)$. The corresponding algebras will be denoted \mathcal{C}_{even} and \mathcal{C}_{odd} .

Now we are ready to consider concrete models of quantum field theory.

3 Massless case

The typical classical continuous equations of motion are first order in time : the free motion (left mover)

$$\dot{p}(x) + p'(x) = 0$$

or KdV equation

$$\dot{p}(x) + p(x)p'(x) + p'''(x) = 0$$


are the typical examples. They are produced by the Poisson structure given above and the Hamiltonians

$$H_0 = \frac{1}{2\gamma} \int p^2(x) dx,$$

$$H_1 = \frac{1}{2\gamma} \int \left(\frac{1}{3} p^3(x) + (p'(x))^2 \right) dx$$

correspondingly. Here we shall treat only the first example. In spite of its triviality it is known to be basic for CFT.

On the rectangular space-time lattice



the simplest generalisation of equation of motion is

$$w_n(t + \Delta) = \tilde{w}_n(t) = w_{n-1}(t).$$

The map (simply a shift)

$$w_n \rightarrow w_{n+1}$$

is evidently the automorphism of algebra \mathcal{A} . The integrability means that we shall be able to construct it explicitly.

We are looking for the operator U such that

$$w_{n+1} = U w_n U^{-1}.$$

Suppose that U is multiplicatively local

$$U = r(w_2) \dots r(w_N).$$

For $n = 1, 2$ or N we get

$$w_{n+1} U = U w_n$$

if

$$w_{n+1} r(w_n) r(w_{n+1}) = r(w_n) r(w_{n+1}) w_n.$$

The basic commutation relations allows us to rewrite this relation as follows

$$r(q^{-2} w_n) w_{n+1} r(w_{n+1}) = r(w_n) w_n r(q^{-2} w_{n+1})$$

and separation of variables leads to the functional equation for the function $r(w)$:

$$w r(w) = c r(q^{-2} w),$$

where c is a constant. We shall normalize this equation as follows

$$\frac{r(qw)}{r(q^{-1}w)} = \frac{1}{w}.$$

Now one can check that the operator U serves as a shift also for the variables w_1, w_2 and w_N if the necessary condition

$$C_1 = C_2$$

is satisfied.

The equation

$$w_1 U = U w_N$$

leads to

$$r(w_1)U = Ur(w_N),$$

so that U can be rewritten as

$$U = r(w_1) \dots r(w_{N-1}).$$

This shows that the expression for U is compatible with the periodicity; it does not depend on a lattice site, we begin the multiplication of $r(w)$ with. The only important things are the order of the factors and that the number of them is equal to $N - 1$ (not N as one could superficially think).

The solution of the functional equation inside the algebra \mathcal{A} (i.e. in power series in w) gives an expression for r as a θ -function

$$r(w) = \sum_{k=-\infty}^{\infty} q^{k^2} w^k,$$

which make sense only if $|q| < 1$. So, the unitary case is nonaccessible for the generic q with values on the unit circle. However, for q being odd root of unity

$$q^{2Q+1} = 1$$

the series

$$r_Q(w) = \sum_{k=-Q}^Q q^{k^2} w^k$$

solves the functional equation.

The solution in the noncompact algebra $\hat{\mathcal{A}}$ exists for all q and is rather simple :

$$\hat{r}(p) = \exp \frac{i}{4\gamma} p^2.$$

One immediately sees that in classical continuous limit the evolution operator U

$$U = e^{-iH\Delta}$$

leads to the classical expression for the Hamiltonian H_0 .

Thus we face the alternative :

1). To work in algebra \mathcal{A} and to bound ourselves in the unitary situation with the q being root of unity. The corresponding dynamical system is finite dimensional, but the dependence on the coupling constant γ is rather erratic.

2). To extend the algebra of observables to $\hat{\mathcal{A}}$. The evolution operator is well defined for all real γ and depends on γ smoothly. However, the algebra of observables is infinite.

More on this alternative will appear soon.

The relevance of our model to CFT becomes explicit in classical continuous case, when one introduces a set of generators of Virasoro algebra

$$S(x) = p^2(x) + p'(x).$$

The Poisson brackets

$$\{S(x), S(y)\} = 2\gamma(S(x) + S(y))\delta'(x - y) + \gamma\delta'''(x - y)$$

known for a long time as a second structure for the KdV equation [11], coincide with those of the Virasoro algebra [12]. The Hamiltonian H_0 can be written now as

$$H_0 = \frac{1}{2\gamma} \int S(x) dx$$

The natural question is if one can introduce a lattice analogue of these formulas. The answer goes as follows [9].

The function $r(w)$ admits the factorization

$$r(w) = S(w)S\left(\frac{1}{w}\right),$$

where $S(w)$ satisfies the following functional equation

$$\frac{S(qw)}{S(q^{-1}w)} = \frac{1}{1+w}.$$

The solution in algebra \mathcal{A} (so that $|q| < 1$) is given by the series

$$S(w) = 1 + \sum_{k=1}^{\infty} q^{\frac{k(k-1)}{2}} \frac{1}{(q^{-1}-q)\dots(q^{-k}-q^k)} w^k,$$

or by the infinite product

$$S(w) = \prod_{k=1}^{\infty} (1 + q^{2k+1}w),$$

or by the expression

$$S(w) = \exp \sum_{k=1}^{\infty} \frac{(-1)^k w^k}{k(q^k - q^{-k})}.$$

This function is well known in combinatorics as a q - Γ -function. The first expression allows also to call it q -exponent, whereas the last one prompts the definition of q -dilogarithm.

In the first guise $S(w)$ satisfies the q -exponential functional equation : let u and v constitute a Weyl pair

$$uv = q^2vu,$$

then

$$S(u)S(v) = S(u+v).$$

I was not able to trace this result to its origine. Apparently it was known in combinatorics long ago. However, its combination with the functional equation leads to the alternative equation

$$S(v)S(u) = S(u+v+q^{-1}uv) = S(u)S(q^{-1}uv)S(v),$$

which apparently was mentioned first in [9]. The first form of it will be used immediately. The second was shown in [13] to be a quantum and q -analogue of the famous pentagon equation for the dilogarithm.

Using the equation

$$S(w_n^{-1})S(w_{n+1}) = S(w_n^{-1} + w_{n+1}q^{-1}w_{n+1}w_n^{-1}),$$

we can easily rewrite the evolution operator U in the form

$$\begin{aligned} U &= S(w_2)S(w_2^{-1})S(w_3)\dots S(w_N)S(w_N^{-1}) = S(w_1^{-1})U(S(w_N^{-1}))^{-1} = \\ &= S(w_1^{-1})S(w_2)S(w_2^{-1})S(w_3)\dots S(w_{N-1}^{-1})S(w_N) = S(s_2)\dots S(s_N), \end{aligned}$$

where

$$s_n = w_{n-1}^{-1} + w_n + q^{-1}w_{n-1}^{-1}w_n.$$

It was shown in [14],[15] that the variables s_n are the natural lattice analogues of the $S(x)$. So, we shall call them the generators of the lattice Virasoro algebra. To avoid missinterpretation let us stress

that there is no q -deformation here, the deformation parameter is only the lattice spacing Δ , which appears in the definition of the classical continuous limit. In particular for $\Delta \rightarrow 0$, $\hbar \rightarrow 0$ we have

$$\frac{1}{4}(1 + s_n) = (1 - \Delta^2 S(x)) + \dots$$

The last expression for the evolution operator U is the analogue of the classical expression for H_0 via $S(x)$.

Now let us turn to the noncompact way. The factorisation

$$\hat{r}(p) = \hat{S}(p)\hat{S}(-p)$$

can be achieved via the solution of the functional equation

$$\frac{\hat{S}(p + \gamma)}{\hat{S}(p - \gamma)} = \frac{1}{1 + e^{ip}}$$

with the solution

$$\hat{S}(p) = \exp \frac{1}{4} \int_{-\infty}^{\infty} \frac{e^{px}}{\sinh(\pi x) \sinh(\gamma x)} \frac{dx}{x},$$

where the singularity of the integral at $x = 0$ is put below the contour of integration. The integral is well defined for $|p| \leq \pi + \gamma$ and then $S(p)$ is continued for all real p by means of the functional equation.

Evaluating the integral through the residues we get the following expression

$$\hat{S}(p) = S_q(w)S_{\hat{q}}^{-1}(\hat{w}),$$

where in the RHS we introduced explicitly dependence of $S(w)$ on q . Whereas each separate factor does not make sense for $|q| = 1$, their ratio does and smoothly depends on γ . We see how two factors, constituting the algebra $\hat{\mathcal{A}}$, harmoniously regularize each other.

It is worthwhile to mention that the pentagon equation in the form

$$\hat{S}(q)\hat{S}(p) = \hat{S}(p)\hat{S}(p+q)\hat{S}(q),$$

where

$$[p, q] = -i\hbar\gamma$$

stays intact in $\hat{\mathcal{A}}$.

Two approaches coincide if we note that the evolution operator U , calculated by the second (noncompact) way, is an outer automorphism of the (compact) algebra \mathcal{A} , reducing to the inner one as soon as it is possible (i.e. for q being root of unity). In this sense the second way is more general.

The real difference appears when we come to the spectral theory. Simple spectrum of algebra \mathcal{A} acquires the multiplicity in $\hat{\mathcal{A}}$. So, the evaluation of spectral characteristics such as partition function depends drastically on the choice of algebra. In particular, in the second way the coupling constants γ and $\frac{\pi^2}{\gamma}$ appear symmetrically. For me it is the origine of all phenomena called duality or mirror symmetry, the appearance of the second quantum group the V. Kac table for minimal models, the duality of the Sine-Gordon and massive Thirring models, etc. This conviction still needs more work for its complete vindication. On this note I shall finish the consideration of the massless example and turn to the massive one.


4 Massive case

The most famous example of the massive model is the Sine-Gordon equation

$$\ddot{\phi}(x, t) - \phi''(x, t) + m^2 \sin \phi = 0$$

for the scalar field $\phi(x, t)$ in two-dimensional space-time. In spite of a vast literature devoted to this model (see references on history and results in [8]), it still continues to reveal its new features. Here I shall show, how the current-like variables can be used to treat the quantum lattice variant of it.


As was already indicated above, the corresponding lattice is to be light-like



with the dynamical variables w_n given on the "initial saw" (the vertical lines symbolize the periodic boundary condition $w_n \equiv w_{n+2M}$).

Consider the equation of motion given as a connection of the dynamical variables

along the elementary plaquette



$$w_N = f(qw_W)w_S(f(qw_E))^{-1} ,$$

where

$$f(w) = \frac{1 + \kappa^2 w}{\kappa^2 + w}$$


and κ^2 is a fixed real positive parameter. Function f maps a unit circle onto itself and is "odd"

$$f\left(\frac{1}{w}\right) = (f(w))^{-1}.$$

The same equation can be rewritten in terms of the dual vertex-like variables ψ

$$\psi_N = \psi_S f\left(q^{-1} \frac{\psi_W}{\psi_E}\right).$$

To check it one is to multiply two last relations according to the picture for two ψ -plaquettes




using the relation like

$$w_W = \psi''_N (\psi'_N)^{-1}$$

and the commutation relations.

To get the classical continuous limit one must make one preliminary step – a change of variables

$$\chi = \begin{cases} \psi \\ \psi^{-1} \end{cases}$$



inverting ψ on each second $SW - NE$ diagonal; consistency with periodicity requires that M is even. The equation in new variables

$$\chi_N = \chi_S^{-1} f(q^{-1} \chi_W \chi_E)$$

turnes into the Sine-Gordon equation in the classical limit if we put

$$\chi = e^{i\phi/2}$$

(no rescaling for $\phi!$), and scale κ^2 in such a way

$$\frac{1}{\Delta^2 \kappa^2} = \frac{m^2}{2}.$$

The form of the lattice S-G equation

$$\sin \frac{1}{4}(\phi_W + \phi_E - \phi_N - \phi_S) = \frac{1}{\kappa^2} \sin \frac{1}{4}(\phi_W + \phi_E + \phi_N + \phi_S)$$

was first introduced in [16] and revisited from the hamiltonian point of view in [17], [10]. However for our goal the form of equation in terms of w (or ψ) is preferable, and we shall stick to it, following [10].

The lattice form of SG equation have the nice properties of its classical continuous contraction, i.e., it can be written as a zero curvature condition, has enough commuting conservation laws, etc.

To show it we introduce the appropriate Lax operator with 2-dimensional auxiliary space. It plays the role of connection along the elementary edge $B \leftarrow A$ of the lattice

$$L_{B \leftarrow A}(\lambda) = \begin{pmatrix} \psi_A^{-1/2} \psi_B^{1/2} & \lambda \psi_A^{-1/2} \psi_B^{-1/2} \\ \lambda \psi_A^{1/2} \psi_B^{1/2} & \psi_A^{1/2} \psi_B^{-1/2} \end{pmatrix}.$$

One sees one more change of normalization $\psi \rightarrow \psi^{1/2}$, but it seems to be inevitable. The equation of motion now takes the form

$$L_{W \leftarrow N}(\kappa^{-1} \lambda) L_{N \leftarrow E}(\kappa \lambda) = L_{W \leftarrow S}(\kappa \lambda) L_{S \leftarrow E}(\kappa^{-1} \lambda)$$

of a zero curvature along the elementary plaquette.

The conservation laws can be obtained by the traditional construction of the transfer matrix; however in our case one is to multiply the Lax operator along the initial saw. With the notation $L_{n-\frac{1}{2}}(\lambda)$ for the Lax operator connecting the vertices with ψ_n and ψ_{n-1} we have

$$t(\lambda) = \text{tr}(L_{2M-\frac{1}{2}}(\kappa \lambda) L_{2M-\frac{3}{2}}(\kappa^{-1} \lambda) \dots L_{\frac{3}{2}}(\kappa \lambda) L_{\frac{1}{2}}(\kappa^{-1} \lambda) C_2^{\frac{1}{2}\sigma_3}),$$

where the martrix $C_2^{\frac{1}{2}\sigma_3}$ takes care of the periodicity. The commutativity

$$[t(\lambda), t(\mu)] = 0$$

is not evident because the Lax operators are not ultralocal; however the ultralocality is restored when one observes that the composed Lax operator

$$\mathbf{L}_n(\lambda) = L_{2n-\frac{1}{2}}(\kappa \lambda) L_{2n-\frac{3}{2}}(\kappa^{-1} \lambda)$$

can be expressed through the canonical pair w_{2n-1}, ψ_{2n-1} (generators of the algebra \mathcal{C}_{odd})

$$\mathbf{L}_n(\lambda) = \lambda \begin{pmatrix} \lambda^{-1} w_{2n-1}^{\frac{1}{2}} + \lambda w_{2n-1}^{-\frac{1}{2}} & \psi_{2n-1}^{\frac{1}{2}} (\kappa w_{2n-1}^{\frac{1}{2}} + \kappa^{-1} w_{2n-1}^{-\frac{1}{2}}) \psi_{2n-1}^{\frac{1}{2}} \\ \psi_{2n-1}^{-\frac{1}{2}} (\kappa^{-1} w_{2n-1}^{\frac{1}{2}} + \kappa w_{2n-1}^{-\frac{1}{2}}) \psi_{2n-1}^{-\frac{1}{2}} & \lambda w_{2n-1}^{\frac{1}{2}} + \lambda^{-1} w_{2n-1}^{-\frac{1}{2}} \end{pmatrix}.$$

The quasiperiodicity of $\mathbf{L}_n(\lambda)$

$$\mathbf{L}_{M+1}(\lambda) = C_2^{\frac{1}{2}\sigma_3} L_1(\lambda) C_2^{-\frac{1}{2}\sigma_3}$$

makes the appearance of $C_2^{\frac{1}{2}\sigma_3}$ in the definition of $t(\lambda)$ quite natural.

The operator $\mathbf{L}_n(\lambda)$ is a variant of the lattice SG Lax operator first introduced in [18]. The possibility of its factorization into the light-cone Lax operator with the vertex-like dynamical variables was realized much later; it was indicated in [19] and stressed in [17].

The Lax operator $\mathbf{L}_n(\lambda)$ is ultralocal and satisfies the ordinary fundamental commutation relation

$$\mathbf{R}\left(\frac{\lambda}{\mu}\right) \mathbf{L}_n(\lambda) \otimes \mathbf{L}_n(\mu) = \mathbf{L}_n(\mu) \otimes \mathbf{L}_n(\lambda) \mathbf{R}\left(\frac{\lambda}{\mu}\right)$$

with the usual 4×4 trigonometric R -matrix. The commutativity of $t(\lambda)$ follows from this. Now $t(\lambda)$ is a polynomial in λ^2 of degree M , and its coefficients are the independent conservation laws in accordance with the number of degrees of freedom.

We can turn to the construction of the evolution operator, which generates the main equation of motion. Some experimentation (or the didactic way, first underlined in [20]) leads to the Ansatz

$$U = \prod r(\kappa^2, w_{\text{even}}) \prod r(\kappa^2, w_{\text{odd}}),$$

where $r(\lambda, w)$ satisfies the functional equation

$$\frac{r(\lambda, qw)}{r(\lambda, q^{-1}w)} = \frac{1 + \lambda w}{\lambda + w}.$$

One can show that the map

$$\psi \rightarrow \tilde{\psi} = U\psi U^{-1}$$

is equivalent to the equation of motion directly by using the commutation relations between ψ -s and w -s and the functional equation. Of course the same is true if one chooses w -s as the main variables. The functional equation is quite inherent to our case. Indeed, it is equivalent to the fundamental commutation relation

$$r\left(\frac{\lambda}{\mu}, w_n\right) L_{n+\frac{1}{2}}(\lambda) L_{n-\frac{1}{2}}(\mu) = L_{n+\frac{1}{2}}(\mu) L_{n-\frac{1}{2}}(\lambda) r\left(\frac{\lambda}{\mu}, w_n\right),$$

which is a variant of the fundamental commutation relation in the quantum space in sense of [3].

It is evident that

$$r(\lambda, w) = \frac{S(w)S(w^{-1})}{S(\lambda w)S(\lambda w^{-1})}$$

is a solution of the functional equation, if $S(w)$ satisfies the functional equation of the previous section. So one faces the alternative of referring to compact or noncompact variant.

In the compact case one can treat the quantum lattice SG equation only for rational coupling constant

$$\gamma = \pi \frac{P}{Q}.$$

The Hilbert space at the given site depends on γ irregularly, its dimension being proportional to Q . In spite of the fact that $S(w)$ makes no sense for $|q| = 1$, the ratio defining $r(\lambda, w)$ can be truncated and leads to the expression for odd Q

$$r(\lambda, w) = 1 + \sum_{k=-\frac{Q-1}{2}}^{\frac{Q-1}{2}} \frac{(1-\lambda)(q-\lambda q^{-1}) \cdots (q^{k-1} - \lambda q^{-k+1})}{(q^{-1} - \lambda q) \cdots (q^{-k} - \lambda q^k)} w^k$$

first found in [21].

In the noncompact case everything depends on γ smoothly, and for $r(\lambda, w)$ we have the expression

$$r(\lambda, w) = \exp \frac{1}{2} \int_{-\infty}^{\infty} \frac{\cosh p\xi (1 - e^{-iw\xi}) d\xi}{\sinh \pi\xi \sinh \gamma\xi \xi},$$

where we put $\lambda = e^w$, which irresistively reminds the Zamolodchikovs phase-shift for the solutions of the SG model [22]. Needless to stress that in our treatment it appears in a completely different function – as a local factor, defining the evolution operator, the variable p (rapidity in the phase-shift interpretation) being a local current-like dynamical variable. The origine and the interpretation of this duality is still unclear but work on this direction is most promising.

Let us repeat, that with respect to dynamics both languages reconcile, if we consider the noncompact U as a outer automorphism of the algebra \mathcal{A} . However, the Hilbert space interpretation of the algebraic formulas is rather different for compact or noncompact choice. This difference seems rather irrelevant in the continuous limit, which is known to exist in quantum case if accompanied with the nontrivial renormalization of the parameter κ .

Now we are in a position to make the comparison of massive and massless case.

5 Discussion and conclusions

In the main body of these lectures I have shown that one can get a rather unified treatment of the massive and massless integrable models on the lattice, using the current-like dynamical variables w . In particular, one gets the possibility to compare both cases on the same common ground.

The massless contraction is rather evident now. The equation of motion

$$w_N = w_W w_S w_E^{-1},$$

which can be obtained from the one in section 4 in the limit $\kappa^2 \rightarrow \infty$ clearly separates: the variables

$$\xi_n = w_{2n} w_{2n+1}^{-1}$$

and

$$\eta_n = w_{2n} w_{2n+1}$$

propagate independently by means of the left or right shift correspondingly. These variables are also current-like

$$\xi_{n+1} \xi_n = q^2 \xi_n \xi_{n+1}$$

$$\eta_n \eta_{n+1} = q^2 \eta_{n+1} \eta_n$$

and mutually commute

$$\xi_n \eta_m = \eta_m \xi_n.$$

The separation in the evolution

$$U_{massive}(w)|_{\kappa^2 \rightarrow \infty} = U_{massless}(\xi) U_{massless}(\eta)$$

is based on the formula

$$r(\kappa^2, w)|_{\kappa^2 = \infty} = r(w)$$

and commutation relations

$$r(w_n) r(w_{n+1}) = r(w_{n+1}) r(q^{-1} w_{n+1}^{-1} w_n)$$

$$r(w_{n+1}) r(w_n) = r(w_n) r(q^{-1} w_n w_{n+1})$$

which follows from the functional equations for $r(w)$ and commutation relations for w_n .

An interesting question appears on this level: what remains of the two lattice Virasoro algebras of massless case in the massive deformation (or more constructively, can they be combined into some new algebraic structure)? This question is not answered yet and deserves more work.

Less superficial results on the interrelation of massive and massless cases appear on the level of the Bethe-Ansatz equations; we cannot discuss them here, as we did not mention Bethe-Ansatz at all. Work in this direction is quite active. In particular the use of Bethe-Ansatz together with the comments on duality in the section 3 must throw a new light on the appearance of traces of two quantum groups in Kac table. The minimal models are to be obtained as massless contraction for special rational γ

$$\gamma = \pi \frac{p}{p+1}.$$

Besides the field-theoretical applications the considerations in these lectures are relevant to some pure mathematical questions. We mention two of them.

1. Quite a few formulas here resemble the treatment of the algebra $U_q(\widehat{sl}(2))$ in [23]. The establishing more close connection is certainly desirable.

2. Classical limit $\hbar \rightarrow 0$ retaining finite lattice site Δ give us a family of models for the integrable symplectic maps. As such it is a valuable addition to the existing results in this field [24], [25].

In these lectures we have treated only the simplest case of the abelian current-like variables. The nonabelian analogue – lattice Kac-Moody algebra – is also known [26] [27]. It is clearly relevant to the lattice WZNW model, which was discussed in [27].

However some principal developments, i.e., the Sugawara construction of Virasoro algebra, are still not finished and work is in progress.

References

- [1] L.D.Faddeev, V.E.Zakharov, *Fun. Anal. and Appl.* **5**, 1972, p.280
- [2] H.Bethe, *Z.Phys.* **71**, 1930, p.205
- [3] L.D.Faddeev, see for example recent lecture notes at ITP, Stony brook, HEP-th April 1994
- [4] R.J.Baxter, *Exactly Solvable Models in Statistical Mechanics*, Academic Press, New-York, 1982
- [5] J.H.Schwarz, *Superstrings*, World Scientific, Singapore, 1986
- [6] A.Belavin, A.Polyakov, A.Zamolodchikov, *Nucl.Phys B* **241**, 1984, p.333
- [7] A.B.Zamolodchikov, *Int.J.Math.Phys.* **A3**, 1988, p.743
- [8] L.D.Faddeev, L.A.Takhtajan, *Hamiltonian Methods in the Theory of Solitons*, Springer, Berlin - New-York, 1987
- [9] L.D.Faddeev, A.Yu.Volkov, *Phys.Lett B* **193**, 1993
- [10] L.D.Faddeev, A.Yu.Volkov, *Lett.Math.Phys.* 1994 (to be published)
- [11] F.Magri, *J.Math.Phys.* **19**, 1978, p.1156
- [12] J.-L.Gervais, *Phys.Lett B* **160**, 1985, p.279
- [13] L.D.Faddeev, R.M.Kashaev, *Mod.Phys.Lett A* **9**, 1994, p.427
- [14] L.D.Faddeev, L.A.Takhtajan, *Springer Lecture Notes, Physics*, **246**, 1986, p.166
- [15] A.Yu.Volkov, *Phys.Lett. A* **167**, 1992, p.345
- [16] R.Hirota, *J.Phys.Soc.Japan*, **43**, 1977, p.2079
- [17] L.D.Faddeev, A.Yu. Volkov, *TMPH* (in russian) **92**, 1992, p.207
- [18] A.G.Izergin, V.E.Korepin, *Lett.Math.Phys.* **5**, 1981, p.199
- [19] V.V.Bazhanov, Yu.G.Stroganov, *J.Stat.Phys.* **51**, 1990, p.799. V.E.Korepin, V.Tarasov, private communication, 1989
- [20] L.D.Faddeev, N.Yu.Reshetikhin, *Ann.Phys.* **167**, 1986, p.227
- [21] V.Fateev, A.B.Zamolodchikov, *Phys.Lett A* **92**, 1982, p.37
- [22] A.B.Zamolodchikov, A.B.Zamolodchikov, *Ann.Phys* **120**, 1979, p.253

- [23] I.B.Frenkel, N.Yu.Reshetikhin, *Commun.Math.Phys.* **146**, 1992, p.1
- [24] J.Moser, A.P.Veselov, *Commun.Math.Phys.* **139**, 1991, p.217
- [25] F.W.Nijhoff et al., *Phys.Rev.* **A 46**, 1992, p.2155; *Phys.Lett.* **A 163**, 1992, p.49
- [26] A.Alexeev, L.Faddeev, M.Semenov-Tian-Shansky, *Commun.Math.Phys.* **149**, 1992, p.335
- [27] L.D.Faddeev, *Quantum Symmetry in Conformal Field Theory by Hamiltonian Methods in New Symmetry Principles in Quantum Field Theory*, NATO ASI Series, Series B: Physics vol. 295, editors J.Fröhlich et al., Plenum Press, New-York and London, 1992